



[www.tasciences.com/clinical-research/](http://www.tasciences.com/clinical-research/)

## Functional Assessment of Pharmacological Telomerase Activators in Human T Cells

Molgora B, Bateman R., Sweeney G., Finger D., Dimler T., Effros R.B., Valenzuela H.F., *Cells Journal*.  
2013 January 14: ISSN 2073—4409

## **Background:**

---

This human study demonstrates that TA-65MD<sup>®</sup> nutritional supplements increase telomerase activity and proliferation in human CD4 and CD8 T cells. It highlights the importance of Telomerase Activation in relation to age-associated pathologies.

## **Study Method:**

---

CD4 and CD8 T cells from healthy subjects were treated with TA-65MD<sup>®</sup>. These samples were measured for telomerase activity 72 hours after primary stimulation. This process was repeated after 18-21 days for a second stimulation.

## **Findings, Method of Action:**

---

TA-65<sup>®</sup> increased telomerase activity in all subjects' T cells during primary and secondary stimulations. The results show that TA-65<sup>®</sup> likely uses the MAPK pathway to activate telomerase. This result was observed in both CD4 and CD8 T cells but most significantly in CD8 T cells that were stimulated for the second time.

## **Findings, Health Effects:**

---

This study confirms that TA-65<sup>®</sup> rapidly induces telomerase activity, which other studies have linked to enhanced healthspan indicators, during an acute viral infection.

The most significant findings of this preliminary study are:

- Clinical situations that require enhanced T cell telomerase activity should benefit from TA-65<sup>®</sup>
- Acute viral infections protocols may benefit from the addition of TA-65<sup>®</sup>

## **Findings, Safety:**

---

No adverse side effects, interactions, or any other negative effects were reported during this study nor have any negative results been reported from any other study associated with TA-65<sup>®</sup>. No unregulated cell growth was observed during this study.